



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Promoting recovery from negative symptoms

**Citation for published version:**

Griffiths, H & McLeod, HJ 2019, Promoting recovery from negative symptoms: An attachment theory perspective. in K Berry, S Bucci & AN Danquah (eds), *Attachment Theory and Psychosis: Current Perspectives and Future Directions*. The International Society for Psychological and Social Approaches to Psychosis Book Series, Routledge, pp. 62-83.

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Attachment Theory and Psychosis

**Publisher Rights Statement:**

This is an Accepted Manuscript of a book chapter published by Routledge in Attachment Theory and Psychosis: Current Perspectives and Future Directions, on 26 Nov 2019, available online:  
<https://www.routledge.com/Attachment-Theory-and-Psychosis-Current-Perspectives-and-Future-Directions/Berry-Bucci-Danquah/p/book/9781138956759>

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## **Promoting recovery from negative symptoms: An attachment theory perspective**

### **ABSTRACT**

Although advances have been made in the understanding and treatment of positive symptoms of psychosis such as paranoid thinking and voice-hearing, negative symptoms such as loss of motivation and reduced social functioning often present a major barrier to recovery. The psychosocial treatment options for negative symptoms should benefit from systematic examination and refinement, both in terms of the models used to understand symptom development and maintenance as well as the specific treatment techniques. We examine the possibility that negative symptom burden is particularly related to insecure-avoidant attachment style and that this manifests in difficulties with understanding one's own mental states and the minds of others. Drawing on key studies that have examined mentalisation and related processes alongside negative symptom profile data, we provide a model of negative symptom formation and maintenance along with a case example that portrays the use of these principles in case formulation and treatment.

Individuals characterised as having the negative symptoms of psychosis participate less in constructive or pleasurable activity and often appear disengaged from interpersonal situations. This group of people also experience less positive emotion, heightened levels of negative affect, and often present as devoid of emotional expression. Their speech can appear impoverished and corresponding mental experiences can be profoundly disrupted. Such difficulties are associated with significant impairments in psychosocial functioning (Hunter & Barry, 2012), lower quality of life (Ho et al, 2014) and poorer recovery (Milev et al, 2005). Response to both pharmacotherapy and psychological interventions appears limited (Fusar-Polis et al, 2014; Velthorst et al, 2015). Furthermore, the diversity of the techniques employed in psychological treatment trials makes it difficult to ascertain potential ingredients that bring about improvement (Thomas, 2015).

To date, many theoretical frameworks underpinning both pharmacological and psychological treatments have mainly reflected the historical understanding of negative symptoms as diminished or absent behaviour, functioning, and experience (Aleman et al., 2016). We argue that, seen through the lens of attachment theory, negative symptoms can be at least partially understood within a developmental framework of adaptation and resilience, as indicators of a learned coping response in the face of overwhelming life adversity that includes both externally experienced events and painful affects that are avoided (Lysaker et al., 2016). Hence, negative symptoms may be seen as responses involving emotional and social withdrawal that emerge from threats to self- security. This alternative perspective may enrich recovery-orientated approaches by supporting attempts to re-engage with life.

## **The Nature of Negative Symptoms**

Even the most severe negative symptoms are known to fluctuate (Ventura et al, 2004). Current cognitive behavioural therapy (CBT) models propose that variations in negative symptoms relate to changes in self-defeating beliefs (Grant & Beck, 2009; Rector et al, 2005). A recent meta-analysis examining changes in negative symptoms in outpatient populations challenges the view that there is limited improvement over time (Savill et al, 2014). Instead, the results suggest that most patients exhibit some recovery of negative symptoms over time. So, rather than viewing negative symptoms as unresponsive to treatment and predictive of an inevitably chronic illness course, the rate of recovery may be helped by developing more refined psychosocial interventions.

Negative symptoms can be observed in individuals at high risk of developing psychosis (Piskulic et al, 2012), they predict likelihood of transition to psychosis (Demjaha et al, 2012, Piskulic et al, 2012) and can feature during first episode presentations (Lyne et al, 2012). Importantly, early difficulties in adjustment and psychosocial functioning link to greater severity of negative symptoms in both first episode and chronic populations (MacBeth & Gumley, 2008). Whilst it has been argued that problems with poor premorbid functioning and the early manifestation of negative symptoms reflect a neurodevelopmental disorder that becomes evident during adolescence (e.g. Fenton & McGlashan, 1991), it has also been established that exposure to socioeconomic deprivation (Drukker et al, 2006), discrimination (Janssen et al, 2003), childhood abuse, neglect and bullying (Varese et al, 2012), and attachment-related disruptions to secure attachments elevate the risk of psychosis. As

Macbeth and Gumley (2008) point out, it would be reasonable to expect that early adversity impacts not only on risk for psychosis, but will also affect premorbid adjustment prior to the development of psychosis. Thus, the difficulty in distinguishing problems in premorbid adjustment, functioning and quality of life from negative symptoms (Malla & Payne, 2005) may be explained within a developmental framework focused on how life experience and learned self-regulatory processes are key risk mechanisms implicated in the early ontogeny of negative symptoms. It has also been proposed that negative symptoms can reflect protective responses to overwhelming life experience including psychosis itself (e.g. Read et al, 2001; Stampfer, 1990). In line with this, psychological adjustment following psychosis, expressed as recovery style, has been shown to both independent and predictive of poorer outcome on a range of variables, including negative symptoms. For example, individuals with a sealing-over recovery style – a way of coping that minimises the significance of symptoms and their impact - had worse negative symptom and quality of life scores at 12 months than those characterised as having integrative or mixed recovery styles (Tait et al, 2003; Thompson et al, 2003).

### **Attachment, mentalization and negative symptoms**

Against this background, the emerging evidence that attachment insecurity is associated with negative symptoms is of considerable interest. The impact of early attachment experience on interpersonal functioning and relationships, emotion regulation, and identity formation is well established. Insecure attachment classifications, specifically dismissing/avoidant attachment, appear to be overrepresented in psychosis (Gumley et al, 2014). Moreover, avoidant attachment, characterised by affective deactivation, interpersonal distancing and difficulty

recalling emotive memories of past experiences, has been specifically associated with negative symptoms in non-clinical and clinical populations (see Table 1). Given that self-report measures may be subject to biases arising from the activation of the attachment system whereby individuals with avoidant attachment styles can tend to report their attachment as secure (e.g. Gumley et al, 2014), these may be underestimates. Recent evidence shows that attachment security, evaluated using the narrative-based Adult Attachment Interview classification system, predicts recovery from negative symptoms at 12 months after controlling for relevant baseline variables (Gumley et al., 2014). There are also particular associations between attachment avoidance and specific negative symptom subtypes such as a loss of the ability to experience pleasure (anhedonia), or social and emotional withdrawal (e.g. Berry et al, 2006, 2007; Wilson & Costanzo, 1996; Korver-Nieberg et al, 2015). In summary, individuals with avoidant attachment style demonstrate restrictions of social-cognitive-affective experience, a pattern frequently observed in people diagnosed with schizophrenia (Gumley, 2011). We propose that these early emotional adaptation processes confer a significant risk for the subsequent emergence of negative symptoms such as asociality, affective flattening, reduced experience of thoughts, diminished motivation, and anhedonia.

Table 1: Attachment Status and Negative Symptoms				
Study	Design	Sample Characteristics	Key Measures	Key findings
i. NON-CLINICAL SAMPLES				
Wilson & Costanzo (1996)	Cross-sectional study of attachment style and sub-clinical psychotic phenomena (schizotypy, attentional impairment, psychosis proneness).	273 university students	Adult Attachment Scale Scale of Attitudes and Experiences (schizotypy)	Insecure-avoidant attachment style associated with negative schizotypy. Secure attachment protects individuals from anhedonia ( $r^2$ =between .06 and .14 for different assessments)
Berry et al. (2006)	Cross-sectional study of attachment style and sub-clinical psychotic phenomena	323 university students	PAM Social Anhedonia Scale	Insecure-avoidant attachment patterns were associated with social anhedonia ( $r=.44$ )
Berry et al (2007)	Cross-sectional analogue study of attachment, negative life events and schizotypy	304 university students	PAM Oxford-Liverpool Inventory of Feelings and Experiences scale (introvertive anhedonia)	Attachment avoidance associated with introvertive anhedonia ( $r=.50$ )
ii. CLINICAL SAMPLE STUDIES				
Ponizovsky et al. (2007)	Cross-sectional with, between and within groups comparisons	30 male inpatients diagnosed with schizophrenia spectrum disorders and 30 age and gender matched controls	Hazan & Shaver questionnaire measure of attachment PANSS	Patients with high PANSS negative symptom scale scores also showed higher rates of insecure-avoidant attachment

Berry et al. (2008)	Prospective study with follow up at 6 months	96 individuals diagnosed with schizophrenia spectrum diagnoses.	PAM PANSS	PANSS negative symptoms were positively correlated with insecure-avoidant attachment at baseline ( $r=.24$ )
Kvrgic et al (2011)	Cross-sectional	127 individuals diagnosed with chronic schizophrenia or schizoaffective disorder	PAM PANSS	No association between attachment avoidance and negative symptoms
Gumley et al. (2014)	12-month longitudinal study with assessments at 6 month intervals	58 first episode psychosis service users in specialist public mental health services	Adult Attachment Interview PANSS	68% of the sample exhibited an insecure attachment style (48.1% dismissing, 20.4% preoccupied). Attachment at baseline predicted negative symptom recovery at 6 and 12 months.
Korver-Nieberg et al (2015)	Cross-sectional study of people diagnosed with schizophrenia spectrum disorders	500 patients	Relationship Questionnaire PANSS	Attachment avoidance was not associated with overall negative symptom scores but there was evidence of associations with social and emotional withdrawal

*Notes*

PAM – Psychosis Assessment Measure; PANSS – Positive and Negative Syndrome Scale



It has also been suggested that negative symptoms may develop as a consequence of reduced capacity for mentalization (Gumley et al 2014; Harder, 2014). Mentalization is conceptually related to other social cognition capacities that have received considerable attention. For example, Frith's (1992) cognitive neurophysiological model of schizophrenia proposed that difficulties with thinking about mental states (Theory of Mind), including monitoring one's own intentional acts, contribute substantially to the development of negative symptoms. The notion that understanding of one's own mind and the mind of others are central to understanding negative symptoms has been corroborated by studies associating Theory of Mind and other social cognition impairments with behavioural indicators of negative symptoms (Sprong et al, 2007), negative/disorganization symptoms (Ventura et al, 2013) and social/functional outcomes (Bora et al, 2009; Fett et al, 2010). It is now known that social cognition interacts with negative self-concepts, such as those related to interpersonal competence, to influence negative symptom expression (Lincoln et al, 2011). In summary, social interaction can be affected by very specific processes (e.g. reading facial expressions) as well as much more complex and integrated processes (e.g. combining multiple sources of information into complex and flexible understandings of the reasons for one's own behaviour and the behaviour of others).

Mentalization and metacognition have at times been used in the psychosis literature more or less interchangeably to refer to the complex representation of mental states of self and others (e.g. Semerari et al, 2003; Lysaker et al 2013). We emphasise mentalization's focus on the way in which these psychological competencies are learnt in attachment or relational contexts. As conceptualised by Peter Fonagy and colleagues, mentalization "describes the way humans make sense of their social world


by imagining the *mental* states (e.g., beliefs, motives, emotions, desires, and needs) that underpin their own and others' behaviors in interpersonal interactions" (Choi-Kain & Gunderson, 2008, p.1127). Mentalization capacity is facilitated through early attachment experience and a caregiver's contingent mirroring of the infant's affective states, and is consequently intrinsically linked to an individual's capacity to employ affect regulation strategies. Secure attachments provide the psychosocial foundations for acquiring the capacity to develop adequate understanding of one's own and others' minds. Effective mentalization allows the individual to organise changing social information in a way that regulates emotions, provides a guide to action, and supports the person's ability to act independently as an agent of their own destiny. When attachment is disrupted, the ability to have complex thoughts about the self and the world can be adversely affected and this results in less resilience when faced with conflicts and dilemmas relating to social interactions or making sense of one's internal experiences (Fonagy & Target, 1997).

Mentalization is more severely compromised in dismissing as opposed to secure or preoccupied attachment states of mind in individuals with psychosis (Macbeth et al, 2011). We identified 14 studies suggesting links between self/other mental state processing and negative symptoms (Buck et al, 2014; Hamm et al, 2012; Lysaker et al, 2005, 2010, 2011, 2015; Macbeth et al, 2011, 2014, 2015; McLeod et al, 2014; Minor & Lysaker, 2014; Mitchell et al, 2012; Nicolo et al, 2012; Rabin et al, 2014). These relationships extend across cognitive, emotional and integrative aspects of mentalizing processes (Macbeth et al, 2015). The majority of studies are cross-sectional, but three indicate that the relationship is linked to the expression of negative symptoms over time in both first episode samples and samples with more

established diagnoses populations (Hamm et al, 2012; Lysaker et al, 2015; McLeod et al, 2014). More refined studies could describe and clarify these relationships in terms of specific metacognitive processes and/or negative symptom subtypes. For example, there is evidence that reduced ability to form and use complex ideas about self and others can exacerbate anhedonia, even in the absence of depressed mood (Buck et al, 2014).

In summary, there is evidence that both avoidant attachment and reduced capacity for mentalization are implicated in negative symptom expression in both early and later stage psychosis. In Figure 1, we propose an explanatory framework, expanding on previous proposals by Gumley et al (2014) and Harder (2014), which suggests how attachment theory could inform our understanding of negative symptoms. In the sections below, we describe potential roles of affect regulation mechanisms and autobiographical memory, considering how these may inform the emergence and maintenance of negative symptoms such as diminished expressivity and avolition/apathy.

**FIG 1: Psychological variables implicated in the development and maintenance of negative symptoms**



## **Affect regulation**

Flat affect, characterised by unchanging facial expression and paucity of gesture, is prominent during first episode and more established presentations (Evensen et al, 2012). However, there is a marked contrast between the emotional experience and expression of individuals diagnosed with schizophrenia, who consistently report less experience of positive affect but greater intensity of negative affect than people included as comparison participants. It is further known that individuals with psychosis have problems with emotion recognition, likely to be present before psychosis onset (Amminger et al, 2012), and linked to social dysfunction, (Kimhy et al, 2012, 2014) and metacognition (Hamm et al, 2012). A recent study of people diagnosed with psychotic disorder demonstrated that this impaired ability to be aware of and to tolerate distressing emotions was associated with increased stress, as measured by both self-report and skin conductance levels (Lincoln et al, 2015). Interestingly, an altered, heightened sensitivity to stress has previously been established as a risk mechanism for psychosis (Myin-Germeys & van Os, 2007). Furthermore, impaired stress tolerance has been associated with negative as well as other symptoms in individuals at high clinical risk of developing psychosis (DeVylder et al, 2013). Although people with psychosis may experience but be less aware of heightened negative affect, they use similar emotion regulation strategies to individuals with anxiety and depressive disorders (Livingstone et al, 2009). However, they use more dysfunctional regulation strategies (e.g. suppression) than non-patient controls (O'Driscoll et al, 2014). To summarise, individuals prone to psychosis demonstrate emotional reactivity in conjunction with a diminished awareness of their own affective state, but use normal, if ultimately maladaptive,

regulatory processes to deal with emotional arousal. This style of dealing with emotional information might shape negative symptom expression.

Adults with dismissing attachment states of mind direct their attention away from emotion and complex mental states, possibly reflecting a developmental adaptation to persistent rejection from caregivers at times of distress. Consistent with this, the degree of critical comments from family members was more strongly associated with negative symptoms than positive symptoms (Barrowclough et al, 2003; for a broader discussion of attachment and expressed emotion see Patterson elsewhere in this volume). At the core of mentalization is ‘mentalized affectivity’ (Fonagy et al, 2002), the capacity to reflect upon the meaning of one’s emotional states as they unfold. The inability to reflect upon emotional distress and its psychosocial causation is not only likely to result in diminished expressivity, but may also block social behaviour, given that distress fails to be communicated. Possible links between unhelpful self-focus (e.g. rumination) and the use of emotional avoidance and withdrawal strategies to deal with interpersonal difficulties have been demonstrated (e.g. McLeod et al, 2014).

The reduced capacity to experience *positive* affect is also likely to play a key role in the development and maintenance of negative symptoms. It is unsurprising that there is little opportunity for positive affect if an individual finds their own or others’ minds confusing and/or threatening and uses emotional, behavioural and social avoidance as primary regulatory strategies. Low expectations of pleasure and success lead to further lost opportunities for meaningful engagement in activity, and become a self-fulfilling prophecy (Rector et al, 2005) that consolidates the maintenance cycle.

Although *anticipation* of pleasure may be impaired, individuals with a high level of

anhedonia can still experience a normal enjoyment response once they are engaged in a pleasurable activity (Gard et al, 2007).

### **Autobiographical memory and personal identity**

The impact of early attachment experience, particularly the ability to recall and reflect on its significance, has long been associated with the development of the sense of self (Cole & Putnam, 1992; Briere, 2002; Conway & Pleydell-Pearce, 2000). The contribution of autobiographical memory to the self is, however, known to be tempered by the need for a coherent and consistent sense of one's self and identity (Conway et al, 2004). For example, when an individual is overwhelmed by life experience, the drive for returning to a stable sense of self identity is given priority over the integration of threatening personal information that goes against one's values or that is inconsistent with one's self-view. In such circumstances, individuals with a dismissing/avoidant attachment style appear to restrict attention to attachment experience, providing sparse personal life-stories in which difficulties appear downplayed. More broadly, this so-called 'overgeneral memory' is considered to be a strategic attempt to control affect associated with specific negative events (Williams et al, 1999). Within various clinical groups, including individuals diagnosed with schizophrenia, lack of specificity in personal memory recall seems to have an adaptive function in the prevention of suicidality (Taylor et al, 2010).

Such self-regulation may come at considerable personal expense. The personal narratives of individuals with dismissing attachment states of mind typically fail to articulate emotional experience, appearing remote and abstract, fragmented, undeveloped and inflexible. Coupled with the ongoing disruption of mentalization,

this distorted recall of key life experience may confer risk for negative symptoms. Attempts to dampen down emotional fluctuations may not only result in restricted expression of affective experience, but also impact on goal-directed activity. In Conway and Pleydell-Pearces's (2000) self-memory system framework, autobiographical memories are seen as forms of mental models that are distinct by virtue of containing self-knowledge and memories for specific episodes experienced in one's life. Importantly, the self-memory system also includes the self's working goals. In this way, autobiographical memory and the sense of oneself as being able to actively influence one's life are intrinsically linked: autobiographical memory 'constrains what the self is, has been and can be' (Conway, 2005). Restricting access to autobiographical memory in order to preserve self-coherence may therefore come about at the expense of engagement in goal-directed activity, and associated development of avolition/apathy.

It has been well documented that people with a diagnosis of schizophrenia have problematic recall of personal past events (Berna et al, 2015). The impact of this may be particularly evident during a crucial period for identity formation i.e. late adolescence (Cuervo-Lombard et al, 2006). Difficulty recalling autobiographical memories or reduced memory specificity is not, of course, specific to schizophrenia, but present in various mental health problems including depression (Liu et al, 2013) and post-traumatic stress (Moore & Zoellner, 2007). Childhood trauma is linked to impairments of memory for earlier life experiences across diagnoses, including in people who have received a diagnosis of schizophrenia (Shannon et al, 2011). Such experiences have been hypothesised to play a specific role in the development of negative symptoms (Read et al, 2001). Relationships have been identified between



negative symptoms and autobiographical memory performance (Aleman et al, 1999; Schmid et al, 2011) and specifically with traumatic memories (McLeod et al, 2006). In a small study exploring the relationships between negative symptoms and autobiographical memory, those with more negative symptoms retrieved fewer specific autobiographical memories and negative symptoms were significantly predicted by both avoidance of traumatic memories and lack of specificity of autobiographical recall (Harrison & Fowler, 2004). Finally, Berna et al (2011) showed that an impaired ability to give meaning to memories that have a strong influence on one's development of self-identity is associated with higher levels of negative symptoms.

Agency can be broadly understood as the basic recognition of the experience of owning and generating one's own thoughts, feelings and behaviours (Lysaker & Leonhardt, 2012). This is a fundamental starting point for experiencing more complex thoughts and is therefore a foundation capacity underpinning metacognition/mentalisation. It is likely that self-regulatory processes associated with impairments of autobiographical memory interact with both the capacity for mentalisation and cognitive variables such as self-efficacy (i.e. appraisals of one's ability to act successfully in the face of challenges). Negative symptoms are associated with reduced self-efficacy (Bentall et al, 2010; Hill & Startup, 2003) as well as the endorsement of attitudes that undermine functioning (e.g. "If you cannot do something well, there is no point in doing it at all") (Ventura et al, 2014). Recent research also suggests that negative symptoms develop in response to self-defeatist beliefs, and these beliefs may affect social functioning because they undermine the ability to see oneself as competent and able (Vaskinn et al, 2015). It seems likely that

relationships between self-efficacy, dysfunctional attitudes and negative symptoms are mutually reinforcing. As an individual's experience and sense of autonomy become ever more restricted, negative symptoms become more entrenched, further restricting opportunities for rich life experience. Given that the sharing of personal stories is fundamental to the development of meaningful human relationships, restricted life narratives may foster increasing isolation. Beliefs about loss, humiliation and entrapment (Iqbal et al, 2000) may then further limit an individual's ability to build an assertive identity as an agentive self who is capable of participation in interpersonal domains.

The costs of these self-regulatory processes are evident in those individuals whose recovery style following the trauma of an acute episode of psychosis has been characterised as 'sealing over'. These individuals tend not to try to integrate or make sense of their psychotic experience in the context of their overall life experience (McGlashan, 1987). Such avoidant coping styles are associated with negative early childhood experience, insecure attachment, negative self-evaluation and insecure identity (Tait et al, 2004). Sealing over is associated with higher levels of negative symptoms (Modestin et al, 2004, 2009; Thompson et al, 2003), poorer social functioning and lower quality of life (McGlashan, 1987; Drayton et al, 1998).

To summarise, it is known that attachment states of mind and the associated capacity for mentalization are significant determinants of the production of self narratives, the evolution of a personal sense of meaning to life experience and the generation of personal identity. Psychological adjustment to life experience that is underpinned by the restriction of attention to personal memory gives priority to the preservation of a coherent personal identity whilst compromising the self's ability to drive future

activity. Such self-regulation may therefore constitute significant risk not only for the emergence of diminished expressivity but also for avolition and motivational difficulties, and impinge significantly on the ongoing evolution of personal identity.

### **Limitations of pure deficit models – critique of contemporary models**

Negative symptoms have typically not been a main treatment target in psychological treatment trials for psychosis over the past two decades (Velthorst et al., 2015). This lack of therapeutic attention is unfortunate given that there is convincing evidence that addressing negative symptoms such as apathy and lack of initiative is a top treatment priority for many people with psychosis (Sterk et al, 2013). Recently more studies have specifically addressed the psychological understanding and treatment of negative symptoms in psychosis with four trials published between 2006-14 where allocation to treatment of a comparison condition was determined at random (see McLeod et al, 2014 for further discussion). The description of the intervention in three of these trials was based on a form of the standard cognitive behavioural therapy model that proposes that low expectations of success in social and functional spheres undermine role engagement and effort expenditure (Grant & Beck, 2009; Rector et al, 2005). This model leads to the use of treatment strategies such as challenging defeatist cognitions, supporting goal setting, testing out beliefs and expectations in real world situations, and making deliberate plans and schedules of activities (Grant et al, 2012). However, the data arising from these trials suggest the need for refinement of the standard CBT model. One indicator of this is that the standard approach may require a large number of treatment sessions to get quite modest symptom improvements (e.g. Grant et al.'s 2012 trial delivered an average of 51 sessions). Also, the standard CBT approach does not seem to be superior to strategies such as

Cognitive Remediation Therapy that do not explicitly focus on changing thoughts and beliefs (Klingberg et al., 2011). These results do suggest that it is possible to make some progress in helping with negative symptoms via psychological methods. We suggest that viewing negative symptoms from the perspective of attachment theory has implications for understanding the delivery of psychological treatments, which may be improved if we develop techniques that more directly target relational factors and the attachment concerns already described. The following case example points to many of these issues.

### **A Case Example**

In the following example, we examine how applying attachment principles to case formulation could lead to useful adaptations of a commonly used therapy, cognitive behavioural therapy. Andreas is a 38-year-old man who lives at home with his elderly mother. His first episode of psychosis at 18 years of age resulted in a hospital admission. At that time he was experiencing distressing hallucinations and paranoia, he was socially withdrawn, had given up attending college and was isolating himself in his bedroom. He was originally encouraged to see his GP by his mother who was increasingly concerned that he was neglecting his self-care and losing weight. When first assessed, he was reluctant to discuss any sources of distress and he denied any need for help or support. The clinical team found it very difficult to engage Andreas in treatment activities. Even gentle encouragement would sometimes result in an agitated and hostile response. This discouraged the clinicians from continuing to push for engagement. A pattern of minimal interaction developed, with the focus of intervention mainly limited to safety monitoring and medication support.

Andreas was eventually referred for psychological treatment. The main areas of concern at that time were his arrested recovery and pervasive failure to engage with social and occupational activities. The assessment and engagement phase of therapy took eight sessions of gradual titration of contact. Andreas' therapeutic goals were to improve his motivation and to understand what was going on in his life, but he was reluctant to set himself any targets in terms of increased activity and was unable to identify potential sources of pleasure. In view of the fact that people with a diagnosis of schizophrenia may experience a form of anhedonia that arises from a fundamental impoverishment of internal experience such that they have substantial difficulty reflecting on and being fully aware of their likes, preferences, wishes, and goals (Buck et al., 2014), the therapist was aware that trying to set goals and engage in activity scheduling in the standard CBT approach may be experienced as stressful rather than therapeutic.

It was therefore agreed to spend the next sessions improving his self-understanding. Although he repeatedly downplayed the severity of his current difficulties, Andreas did slowly open up about early difficulties attributable to his mother's alcoholism. He had been repeatedly sent to live with relatives at times when his mother was unable to cope. During many of these episodes he was sexually and physically abused. Although he made these disclosures, Andreas continued to describe many of these experiences in a dispassionate and unemotional way during this early phase of therapy. Although slow exploration of early memories and perceptions of key experiences were summarised by the therapist into a time-line, Andreas found it extremely difficult to reflect on how his early experiences might have had an impact on his current situation. Throughout therapy, the therapist maintained an awareness

that people exhibiting an avoidant attachment style will likely experience the therapeutic relationship to be a source of stress and threat which may elicit deactivating emotion regulation strategies such as social withdrawal and down-regulation of metacognition (Gumley et al, 2014; Lysaker et al., 2015). This prompted the therapist to be particularly sensitive to relational patterns during sessions, attending to pacing and timing of sessions, continuously assessing the client's tolerance for addressing previously avoided emotional material, and actively working to strengthen the security of the therapeutic relationship. The experienced therapist required regular supervision to support this kind of work. Such strategies facilitated the development of a therapeutic relationship that the therapist believed was able to provide a helpful emotional experience (Daly & Mallinckrodt, 2009; Mallinckrodt & Jeong, 2015).

Adding these attachment dimensions to Andreas' therapeutic formulation also guided the matching of therapeutic strategies to the level of metacognitive capacity exhibited by him. The formulation that emerged over time was that Andreas had coped with repeated episodes of abusive behaviour from caregivers through social and emotional withdrawal. His aim was to become as inconspicuous as possible. Andreas' mental model of others became dominated by the view that people were at best unreliable as sources of support and at worst, dangerous and abusive. Unsurprisingly, he also seemed to have an impoverished capacity to think about the reasons why others act in the way that they do. Having had little opportunity to learn effective self-regulation, attempts to consider his own mental states were similarly experienced as overwhelmingly painful.

In this case we can see how there were barriers to implementing a standard CBT approach. Andreas' difficulties with engagement with professional carers and the limited emotional connectedness to his mother both reflect a fundamental disruption of the ability to use relationships to meet challenges, soothe negative affect or provide a source of encouragement and drive to engage with life challenges. Goals that involved any level of social connectedness and interpersonal relating activated distress, confusion and withdrawal behaviours. The goal setting process was further impeded by Andreas' diminished ability to reflect on his autobiographical memories and lack of access to previous sources of reward and meaning. Engaging in standard CBT techniques of exploring thoughts and feelings about others (e.g. expectations of low social acceptance) was difficult for Andreas because he experienced them as highly aversive.

Instead, Andreas' therapist used the formulation to guide treatment in the following ways. First, the carefully titrated engagement and assessment process created an opportunity to decrease the threat associated with interpersonal contact. The next therapeutic task was to help Andreas explore and differentiate his thoughts, feelings, goals, aspirations and wishes (e.g. see Lysaker et al., 2011 for a description of principles underpinning this approach). This then led to the therapist providing a model of her mind by describing in transparent terms some of the thoughts and understandings that occurred as therapy progressed. At a more subtle level, the therapy interaction also provided opportunities to shape up expressive behaviours that facilitated social interactions (e.g. the therapist explicitly provided feedback when she noticed an expression of affect passing across Andreas' face). This provided an opportunity for Andreas to practise noticing and reflecting on mental states in a way

that minimised threat, ambiguity, and misinterpretation. He was gradually encouraged to communicate to the therapist when he was experiencing high levels of affect during the session. The therapist then prompted the use of strategies such as grounding and soothing techniques. Once the basic ability to engage in conversations about thoughts, feelings, goals, and aspirations had been re-established, therapy could move on to more conventional CBT strategies such as setting between session behavioural tasks, examining the accuracy of thoughts and beliefs that undermine behaviour change (e.g. engaging socially with others will always be unrewarding), and gradually expanding the range of behaviours that Andreas was willing to attempt (e.g. scheduling an achievable array of master and pleasure focused tasks).

This approach uses an understanding of attachment states of mind and the associated problems with mentalisation to reach people who are less responsive to standard CBT for negative symptom strategies. Our suggestions share commonalities with other variants of mentalization-based therapy which have recently been adapted for individuals diagnosed with psychotic disorder to facilitate therapeutic alliance (Brent, 2015) and social functioning (Weijers et al, 2016). Most importantly, these approaches all identify the need to specifically target disturbances in thinking capacities related to awareness of self and others. Sensitively titrated therapy adapted along mentalization-based therapy principles provides an opportunity to learn the interpersonal thinking and relating skills needed to re-engage with more traditional goal and task focused CBT.

## **Conclusions**



There is increasing recognition that interventions for negative symptoms of schizophrenia require more attention. In fact, data collected from people diagnosed with psychoses suggest that improving motivation and self understanding are highly ranked treatment targets (Byrne & Morrison, 2015; Sterk et al , 2013). In this chapter, we have tried to articulate an important but often overlooked issue for psychological treatments: there is a need to tailor the treatment strategies not just to address cognitive maintenance factors (e.g. self-defeating cognitions) but also to take account of relevant sources of resilience (Dudley et al, 2011) and interactions between distal risk factors such as abuse, neglect, and disrupted attachment (Nolen-Hoeksema & Watkins, 2011). For some people exhibiting negative symptoms, attempts at standard approaches (e.g. CBT, social skills training) may not be possible until the impact of disrupted attachment has been assessed, understood, and incorporated into the formulation and treatment plan. As our case study portrays, without attention being paid to these issues the likelihood of successful recovery is greatly reduced.

## References

Aleman, A., Hijman, R., de Haan, E. H., & Kahn, R. S. (1999). Memory impairment in schizophrenia: a meta-analysis. *American Journal of Psychiatry*, 156(9), 1358-1366.

Aleman, A., Lincoln, T. M., Bruggeman, R., Melle, I., Arends, J., Arango, C., & Knegtering, H. (2016). Treatment of negative symptoms: Where do we stand, and where do we go? *Schizophrenia Research*.  
<http://doi.org/10.1016/j.schres.2016.05.015>

Amminger, G. P., Schäfer, M. R., Papageorgiou, K., Klier, C. M., Schlögelhofer, M., Mossaheb, N., ... & McGorry, P. D. (2012). Emotion recognition in individuals at clinical high-risk for schizophrenia. *Schizophrenia Bulletin*, 38(5), 1030-1039.

Barrowclough, C., Tarrier, N., Humphreys, L., Ward, J., Gregg, L., & Andrews, B. (2003). Self-esteem in schizophrenia: relationships between self-evaluation, family attitudes, and symptomatology. *Journal of Abnormal Psychology*, 112(1), 92.

Bentall, R. P., Simpson, P. W., Lee, D. A., Williams, S., Elves, S., Brabbins, C., & Morrison, A. P. (2010). Motivation and avolition in schizophrenia patients: The role of self-efficacy. *Psychosis*, 2(1), 12-22.

Berna, F., Potheegadoo, J., Aouadi, I., Ricarte, J. J., Allé, M. C., Coutelle, R., ... & Danion, J. M. (2015). A Meta-Analysis of Autobiographical Memory Studies in Schizophrenia Spectrum Disorder. *Schizophrenia Bulletin*, sbv099.

Berna, F., Bennouna-Greene, M., Potheegadoo, J., Verry, P., Conway, M. A., & Danion, J. M. (2011). Impaired ability to give a meaning to personally significant events in patients with schizophrenia. *Consciousness and Cognition*, 20(3), 703-711.

Berry, K., Band, R., Corcoran, R., Barrowclough, C., & Wearden, A. (2007). Attachment styles, earlier interpersonal relationships and schizotypy in a non-clinical sample. *Psychology and Psychotherapy: Theory, Research and Practice*, 80(4), 563-576.

Berry, K., Barrowclough, C., & Wearden, A. (2008). Attachment theory: A framework for understanding symptoms and interpersonal relationships in psychosis. *Behaviour Research and Therapy*, 46, 1275-1282

Berry, K., Wearden, A., Barrowclough, C., & Liversidge, T. (2006). Attachment styles, interpersonal relationships and psychotic phenomena in a non-clinical student sample. *Personality and Individual Differences*, 41(4), 707-718.

Bora, Emre, Yucel, Murat, & Pantelis, Christos. (2009). Theory of mind impairment in schizophrenia: Meta-analysis. *Schizophrenia Research*, 109(1), 1-9.

Brent, B. K. (2015). A Mentalization-Based Approach to the Development of the Therapeutic Alliance in the Treatment of Schizophrenia. *Journal of Clinical Psychology*, 71(2), 146-156.

Briere, J. (2002). Treating Adult Survivors of Severe Childhood Abuse. *The APSAC handbook on child maltreatment*, 175.

Buck, K. D., McLeod, H. J., Gumley, A., Dimaggio, G., Buck, B. E., Minor, K. S., ... & Lysaker, P. H. (2014). Anhedonia in prolonged schizophrenia spectrum patients with relatively lower vs. higher levels of depression disorders: Associations with deficits in social cognition and metacognition. *Consciousness and Cognition*, 29, 68-75.

Byrne, R., & Morrison, A. P. (2014). Service users' priorities and preferences for treatment of psychosis: a user-led Delphi study. *Psychiatric Services*.

Choi-Kain, L. W., & Gunderson, J. G. (2008). Mentalization: ontogeny, assessment, and application in the treatment of borderline personality disorder. *The American journal of psychiatry*, 165(9), 1127-1135.

Cole, P. M., & Putnam, F. W. (1992). Effect of incest on self and social functioning: A developmental psychopathology perspective. *Journal of consulting and clinical psychology*, 60(2), 174.

Conway, M. A., & Pleydell-Pearce, C. W. (2000). The construction of autobiographical memories in the self-memory system. *Psychological review*, 107(2), 261.

Conway, M. A., Singer, J. A. & Tagini, A. (2004) The Self and Autobiographical Memory: Correspondence and Coherence. *Social Cognition*, 22(5), 491-529.

Conway, Martin A. (2005). Memory and the Self. *Journal of Memory and Language*, 53(4), 594-628.

Cuervo-Lombard, C., Jovenin, N., Hedelin, G., Rizzo-Peter, L., Conway, M., & Danion, J. (2007). Autobiographical memory of adolescence and early adulthood events: An investigation in schizophrenia. *Journal of the International Neuropsychological Society*, 13(2), 335-343.

Daly, K. D., & Mallinckrodt, B. (2009). Experienced therapists' approach to psychotherapy for adults with attachment avoidance or attachment anxiety. *Journal of Counseling Psychology*, 56(4), 549–563. <http://doi.org/10.1037/a0016695>

Devylder, J. E., Ben-David, S., Schobel, S. A., Kimhy, D., Malaspina, D., & Corcoran, C. M. (2013). Temporal association of stress sensitivity and symptoms in individuals at clinical high risk for psychosis. *Psychological medicine*, 43(02), 259-268.

Demjaha, A., Valmaggia, L., Stahl, D., Byrne, M., & McGuire, P. (2012). Disorganization/cognitive and negative symptom dimensions in the at-risk mental state predict subsequent transition to psychosis. *Schizophrenia bulletin*, 38(2), 351-359.

Drayton, M., Birchwood, M., & Trower, P. (1998). Early attachment experience and recovery from psychosis. *British Journal of Clinical Psychology*, 37(3), 269-284.

Drukker M, Krabbendam L, Driessen G, Van Os J. (2006) Social disadvantage and schizophrenia. A combined neighbourhood and individual-level analysis. *Soc Psychiatry Psychiatr Epidemiol*;41:595–604. <sup>[1]</sup><sub>SEP</sub>

Dudley, R., Kuyken, W., & Padesky, C. A. (2011). Disorder specific and trans-diagnostic case conceptualisation. *Clinical Psychology Review*, 31(2), 213-224.

Evensen, J., Røssberg, J. I., Barder, H., Haahr, U., ten Velden Hegelstad, W., Joa, I., ... & Friis, S. (2012). Flat affect and social functioning: a 10year follow-up study of first episode psychosis patients. *Schizophrenia research*, 139(1), 99-104.

Fenton, W. S., & McGlashan, T. H. (1991). Natural history of schizophrenia subtypes: II. Positive and negative symptoms and long-term course. *Archives of general psychiatry*, 48(11), 978-986.

Fett, Anne-Kathrin J., Viechtbauer, Wolfgang, Dominguez, Maria-de-Gracia, Penn, David L., Van Os, Jim, & Krabbendam, Lydia. (2010). The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: A meta-analysis. *Neuroscience and Biobehavioral Reviews*, 35(3), 573-588.

Fonagy, P., Gergely, G., Jurist, E., et al ( 2002) *Affect Regulation, Mentalization and the Development of the Self*. New York: Other Press.

Fonagy, P., & Target, M. (1997). Attachment and reflective function: Their role in self-organization. *Development and Psychopathology*, 9(4), 679-700.

Frith, C. (1992). *The cognitive neuropsychology of schizophrenia* (Essays in cognitive psychology). Hove: L. Erlbaum.

Fusar-Poli, P., Papanastasiou, E., Stahl, D., Rocchetti, M., Carpenter, W., Shergill, S., & McGuire, P. (2014). Treatments of negative symptoms in schizophrenia: meta-analysis of 168 randomized placebo-controlled trials. *Schizophrenia bulletin*, sbu170.

Gard, D. E., Kring, A. M., Gard, M. G., Horan, W. P., & Green, M. F. (2007). Anhedonia in schizophrenia: distinctions between anticipatory and consummatory pleasure. *Schizophrenia research*, 93(1), 253-260.

Grant, P. M., & Beck, A. T. (2009). Defeatist beliefs as a mediator of cognitive impairment, negative symptoms, and functioning in schizophrenia. *Schizophrenia bulletin*, 35(4), 798-806.

Grant, P. M., Huh, G. A., Perivoliotis, D., Stolar, N. M., & Beck, A. T. (2012). Randomized trial to evaluate the efficacy of cognitive therapy for low-functioning patients with schizophrenia. *Archives of General Psychiatry*, 69(2), 121-127.

Gumley, A. (2011). Metacognition, affect regulation and symptom expression: A transdiagnostic perspective. *Psychiatry research*, 190(1), 72-78.

Gumley, A. I., Schwannauer, M., Macbeth, A., Fisher, R., Clark, S., Rattrie, L., ... & Birchwood, M. (2014). Insight, duration of untreated psychosis and attachment in first-episode psychosis: prospective study of psychiatric recovery over 12-month follow-up. *The British Journal of Psychiatry*, 205(1), 60-67.

Gumley, A. I., Taylor, H. E. F., Schwannauer, M., & MacBeth, A. (2014). A systematic review of attachment and psychosis: measurement, construct validity and outcomes. *Acta Psychiatrica Scandinavica*, 129(4), 257-274.

Hamm, J. A., Renard, S. B., Fogley, R. L., Leonhardt, B. L., Dimaggio, G., Buck, K. D., & Lysaker, P. H. (2012). Metacognition and social cognition in schizophrenia: stability and relationship to concurrent and prospective symptom assessments. *Journal of clinical psychology*, 68(12), 1303-1312.

Harder, S. (2014). Attachment in Schizophrenia—implications for research, prevention, and Treatment. *Schizophrenia bulletin*, 40(6), 1189-1193.

Harrison, C. L., & Fowler, D. (2004). Negative symptoms, trauma, and autobiographical memory: an investigation of individuals recovering from psychosis. *The Journal of nervous and mental disease*, 192(11), 745-753.

Hill, Kimberley, & Startup, Mike. (2012). The relationship between internalized stigma, negative symptoms and social functioning in schizophrenia: The mediating role of self-efficacy. *Psychiatry Research*, Psychiatry Research.



Ho, B. C., Nopoulos, P., Flaum, M., Arndt, S., & Andreasen, N. C. (2014). Two-year outcome in first-episode schizophrenia: predictive value of symptoms for quality of life. *Focus*.

Hunter, R., & Barry, S. (2012). Negative symptoms and psychosocial functioning in schizophrenia: neglected but important targets for treatment. *European Psychiatry*, 27(6), 432-436.

Iqbal, Birchwood, Chadwick, & Trower. (2000). Cognitive approach to depression and suicidal thinking in psychosis. 2. Testing the validity of a social ranking model. *The British Journal of Psychiatry : The Journal of Mental Science*, 177, 522-8.

Janssen, I., Hanssen, M., Bak, M., Bijl, R. V., De Graaf, R., Vollebergh, W., ... & Van Os, J. (2003). Discrimination and delusional ideation. *The British Journal of Psychiatry*, 182(1), 71-76.

Kimhy, D., Vakhrusheva, J., Jobson-Ahmed, L., Tarrier, N., Malaspina, D., & Gross, J. J. (2012). Emotion awareness and regulation in individuals with schizophrenia: Implications for social functioning. *Psychiatry research*, 200(2), 193-201.

Kimhy, D., Vakhrusheva, J., Khan, S., Chang, R. W., Hansen, M. C., Ballon, J. S., ... & Gross, J. J. (2014). Emotional granularity and social functioning in individuals with schizophrenia: An experience sampling study. *Journal of psychiatric research*, 53, 141-148.

Klingberg, S., Wölwer, W., Engel, C., Wittorf, A., Herrlich, J., Meisner, C., ... & Wiedemann, G. (2011). Negative symptoms of schizophrenia as primary target of cognitive behavioral therapy: results of the randomized clinical TONES study. *Schizophrenia bulletin*, 37(suppl 2), S98-S110.

Korver-Nieberg, Berry, Meijer, De Haan, & Ponizovsky. (2015). Associations between attachment and psychopathology dimensions in a large sample of patients with psychosis. *Psychiatry Research*, 228(1), 83-88.

Kvrgic, S., Beck, E., Cavelti, M., Kossowsky, J., Stieglitz, R., & Vauth, R. (2011). Focusing on the adult attachment style in schizophrenia in community mental health centres: Validation of the Psychosis Attachment Measure (PAM) in a German speaking sample. *International Journal of Social Psychiatry*, 58, 362–373.

Lincoln, T., Mehl, S., Kesting, M., & Rief, W. (2011). Negative Symptoms and Social Cognition: Identifying Targets for Psychological Interventions. *Schizophrenia Bulletin*, 37(Suppl2), S23-S32.

Lincoln, T. M., Hartmann, M., Köther, U., & Moritz, S. (2015). Dealing with feeling: Specific emotion regulation skills predict responses to stress in psychosis. *Psychiatry research*.

Livingstone, K., Harper, S., & Gillanders, D. (2009). An exploration of emotion regulation in psychosis. *Clinical psychology & psychotherapy*, 16(5), 418-430.

Liu, X., Li, L., Xiao, J., Yang, J. and Jiang, X. (2013), Abnormalities of autobiographical memory of patients with depressive disorders: A meta-analysis. *Psychology and Psychotherapy: Theo, Res, Pra*, 86: 353–373. doi: 10.1111/j.2044-8341.2012.02077.x

Lyne, J., O'Donoghue, B., Owens, E., Renwick, L., Madigan, K., Kinsella, A., ... & O'Callaghan, E. (2012). Prevalence of item level negative symptoms in first episode psychosis diagnoses. *Schizophrenia research*, 135(1), 128-133.

Lysaker, P. H., & Leonhardt, B. L. (2012). Agency: its nature and role in recovery from severe mental illness. *World Psychiatry*, 11(3), 165-166.

Lysaker, P. H., Buck, K. D., Carcione, A., Procacci, M., Salvatore, G., Nicolò, G., & Dimaggio, G. (2011). Addressing metacognitive capacity for self reflection in the psychotherapy for schizophrenia: A conceptual model of the key tasks and processes. *Psychology and Psychotherapy: Theory, Research and Practice*, 84, 58–69.  
<http://doi.org/10.1348/147608310X520436>

Lysaker, P. H., Carcione, A., Dimaggio, G., Johannesen, J. K., Nicolò, G., Procacci, M., & Semerari, A. (2005). Metacognition amidst narratives of self and illness in schizophrenia: associations with neurocognition, symptoms, insight and quality of life. *Acta Psychiatrica Scandinavica*, 112(1), 64-71.

Lysaker, P. H., Gumley, A., Brüne, M., Vanheule, S., Buck, K. D., & Dimaggio, G. (2011). Deficits in the ability to recognize one's own affects and those of others: associations with neurocognition, symptoms and sexual trauma among persons with schizophrenia spectrum disorders. *Consciousness and Cognition*, 20(4), 1183-1192.

Lysaker, P. H., Kukla, M., Belanger, E., White, D. A., Buck, K. D., Luther, L., et al. (2016). Individual Psychotherapy and Changes in Self-Experience in Schizophrenia: A Qualitative Comparison of Patients in Metacognitively Focused and Supportive Psychotherapy. *Psychiatry: Interpersonal and Biological Processes*, 78(4), 305–316. <http://doi.org/10.1080/00332747.2015.1063916>

Lysaker, P. H., Kukla, M., Dubreucq, J., Gumley, A., McLeod, H., Vohs, J. L., et al. (2015). Metacognitive deficits predict future levels of negative symptoms in schizophrenia controlling for neurocognition, affect recognition, and self-expectation of goal attainment. *Schizophrenia Research*. <http://doi.org/10.1016/j.schres.2015.06.015>

Lysaker, P. H., Shea, A. M., Buck, K. D., Dimaggio, G., Nicolò, G., Procacci, M., ... & Rand, K. L. (2010). Metacognition as a mediator of the effects of impairments in neurocognition on social function in schizophrenia spectrum disorders. *Acta Psychiatrica Scandinavica*, 122(5), 405-413.

MacBeth, A., & Gumley, A. (2008). Premorbid adjustment, symptom development and quality of life in first episode psychosis: a systematic review and critical reappraisal. *Acta Psychiatrica Scandinavica*, 117(2), 85-99.

MacBeth, A., Gumley, A., Schwannauer, M., Carcione, A., Fisher, R., McLeod, H. J., & Dimaggio, G. (2014). Metacognition, symptoms and premorbid functioning in a First Episode Psychosis sample. *Comprehensive Psychiatry*, 55(2), 268-273.

MacBeth, A., Gumley, A., Schwannauer, M., Carcione, A., McLeod, H. J., & Dimaggio, G. (2015). Metacognition in First Episode Psychosis: Item Level Analysis of Associations with Symptoms and Engagement. *Clinical Psychology & Psychotherapy*.

MacBeth, A., Gumley, A., Schwannauer, M., & Fisher, R. (2011). Attachment states of mind, mentalization, and their correlates in a first-episode psychosis sample. *Psychology and Psychotherapy: Theory, Research and Practice*, 84(1), 42-57.

Malla, A., & Payne, J. (2005). First-episode psychosis: psychopathology, quality of life, and functional outcome. *Schizophrenia Bulletin*, 31(3), 650-671.

Mallinckrodt, B., & Jeong, J. (2015). Meta-analysis of client attachment to therapist: Associations with working alliance and client pretherapy attachment. *Psychotherapy*, 52(1), 134-139.

Mcglashan, T. H. (1987). Recovery Style from Mental Illness and Long-Term Outcome. *The Journal of Nervous and Mental Disease*, 175(11), 681-685.

McLeod, H. J., Wood, N., & Brewin, C. R. (2006). Autobiographical memory deficits in schizophrenia. *Cognition & Emotion*, 20(3-4), 536-547.

McLeod, H. J., Gumley, A. I., MacBeth, A., Schwannauer, M., & Lysaker, P. H. (2014). Metacognitive functioning predicts positive and negative symptoms over 12 months in first episode psychosis. *Journal of psychiatric research*, 54, 109-115.

McLeod, H. J., Gumley, A., & Schwannauer, M. (2014). The Impact of Metacognition on the Development and Maintenance of Negative Symptoms. *Social Cognition and Metacognition in Schizophrenia: Psychopathology and Treatment Approaches*, 115.

Milev, P., Ho, B. C., Arndt, S., & Andreasen, N. C. (2005). Predictive values of neurocognition and negative symptoms on functional outcome in schizophrenia: a longitudinal first-episode study with 7-year follow-up. *American Journal of Psychiatry*, 162(3), 495-506.

Minor, K. S., & Lysaker, P. H. (2014). Necessary, but not sufficient: Links between neurocognition, social cognition, and metacognition in schizophrenia are moderated by disorganized symptoms. *Schizophrenia research*, 159(1), 198-204.

Mitchell, L. J., Gumley, A., Reilly, E. S., Macbeth, A., Lysaker, P., Carcione, A., & Dimaggio, G. (2012). Metacognition in forensic patients with schizophrenia and a past history of interpersonal violence: an exploratory study. *Psychosis*, 4(1), 42-51.

Modestin J, Caveng I, Wehrli M, Malti T (2009) Correlates of coping styles in psychotic illness - an extension study. *Psychiatr Res*;168:50-56.

- Moore, S. A., & Zoellner, L. A. (2007). Overgeneral autobiographical memory and traumatic events: an evaluative review. *Psychological bulletin*, 133(3), 419.
- Myin-Germeys, I., & van Os, J. (2007). Stress-reactivity in psychosis: evidence for an affective pathway to psychosis. *Clinical psychology review*, 27(4), 409-424.
- Nicolo, G., Dimaggio, G., Popolo, R., Carcione, A., Procacci, M., Hamm, J., ... & Lysaker, P. H. (2012). Associations of metacognition with symptoms, insight, and neurocognition in clinically stable outpatients with schizophrenia. *The Journal of nervous and mental disease*, 200(7), 644-647.
- Nolen-Hoeksema, S., & Watkins, E. R. (2011). A heuristic for developing transdiagnostic models of psychopathology explaining multifinality and divergent trajectories. *Perspectives on Psychological Science*, 6(6), 589-609.
- O'Driscoll, C., Laing, J., & Mason, O. (2014). Cognitive emotion regulation strategies, alexithymia and dissociation in schizophrenia, a review and meta-analysis. *Clinical psychology review*, 34(6), 482-495.
- Patterson, P. (2013). Attachment, loss and expressed emotion. *Psychosis and emotion: The role of emotions in understanding psychosis, therapy and recovery*, 136-148.

- Piskulic, D., Addington, J., Cadenhead, K. S., Cannon, T. D., Cornblatt, B. A., Heinssen, R., ... & McGlashan, T. H. (2012). Negative symptoms in individuals at clinical high risk of psychosis. *Psychiatry research*, 196(2), 220-224.
- Ponizovsky, A. M., Nechamkin, Y., & Rosca, P. (2007). Attachment patterns are associated with symptomatology and course of schizophrenia in male inpatients. *American Journal of Orthopsychiatry*, 77(2), 324.
- Rabin, S., Hasson-Ohayon, I., Avidan, M., Rozencwaig, S., Shalev, H., & Kravetz, S. (2014). Metacognition in Schizophrenia and Schizotypy: Relation to Symptoms of Schizophrenia, Traits of Schizotypy and Social Quality of Life. *Israel Journal Of Psychiatry And Related Sciences*, 51(1), 44-53.
- Read, J., Perry, B. D., Moskowitz, A., & Connolly, J. (2001). The contribution of early traumatic events to schizophrenia in some patients: a traumagenic neurodevelopmental model. *Psychiatry*, 64(4), 319-345.
- Rector, N. A., Beck, A. T., & Stolar, N. (2005). The negative symptoms of schizophrenia: a cognitive perspective. *Canadian Journal of Psychiatry*, 50(5), 247-257.
- Savill, M., Banks, C., Khanom, H., & Priebe, S. (2015). Do negative symptoms of schizophrenia change over time? A meta-analysis of longitudinal data. *Psychological medicine*, 45(08), 1613-1627.



- Schmid, L. A., Lässer, M. M., Herold, C. J., Seidl, U., & Schröder, J. (2010). Autobiographical memory deficits and their relationship with symptom dimensions in older patients with chronic schizophrenia. *Schizophrenia Research*, 117(2), 216.
- Shannon, C., Douse, K., McCusker, C., Feeney, L., Barrett, S., Mulholland, C., 2011. The association between childhood trauma and memory functioning in schizophrenia. *Schizophrenia Bulletin* 37 (3), 531–537.
- Sprong, M., Schothorst, P., Vos, E., Hox, J., & Van Engeland, H. (2007). Theory of mind in schizophrenia: Meta-analysis. *The British Journal of Psychiatry : The Journal of Mental Science*, 191, 5-13.
- Stampfer, H. G. (1990). "Negative Symptoms": A Cu Mu Latlve Trauma Stress Disorder?. *Australian and New Zealand Journal of Psychiatry*, 24(4), 516-528.
- Sterk, B., Winter, R. I., Muis, M., & de Haan, L. (2013). Priorities, satisfaction and treatment goals in psychosis patients: an online consumer's survey. *Pharmacopsychiatry*. <http://doi.org/10.1055/s-0032-1>
- Tait, L., Birchwood, M., & Trower, P. (2003). Predicting engagement with services for psychosis: insight, symptoms and recovery style. *The British Journal of Psychiatry*, 182(2), 123-128.
- Tait, L., Birchwood, M., & Trower, P. (2004). Adapting to the challenge of psychosis: Personal resilience and the use of sealing-over (avoidant) coping

strategies. *The British Journal of Psychiatry : The Journal of Mental Science*, 185, 410-5.

Taylor, Peter James, et al. "Memory specificity as a risk factor for suicidality in non-affective psychosis: The ability to recall specific autobiographical memories is related to greater suicidality." *Behaviour Research and Therapy* 48.10 (2010): 1047-1052.

Thomas, N. (2015). What's really wrong with cognitive behavioral therapy for psychosis? *Frontiers in Psychology*, 6. <http://doi.org/10.3389/fpsyg.2015.00323>

Thompson, K. N., McGorry, P. D., & Harrigan, S. M. (2003). Recovery style and outcome in first-episode psychosis. *Schizophrenia Research*, 62(1), 31-36.

Varese, F., Smeets, F., Drukker, M., Lieverse, R., Lataster, T., Viechtbauer, W., ... & Bentall, R. P. (2012). Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective-and cross-sectional cohort studies. *Schizophrenia Bulletin*, 38(4), 661-671.

Vaskinn, Ventura, Andreassen, Melle, & Sundet. (2015). A social path to functioning in schizophrenia: From social self-efficacy through negative symptoms to social functional capacity. *Psychiatry Research*,.

Velthorst, E., Koeter, M., van der Gaag, M., Nieman, D. H., Fett, A. K., Smit, F., ... & de Haan, L. (2015). Adapted cognitive-behavioural therapy required for targeting negative symptoms in schizophrenia: meta-analysis and meta-regression. *Psychological medicine*, 45(03), 453-465.

Ventura, J., Nuechterlein, K. H., Green, M. F., Horan, W. P., Subotnik, K. L., & Mintz, J. (2004). The timing of negative symptom exacerbations in relationship to positive symptom exacerbations in the early course of schizophrenia. *Schizophrenia research*, 69(2), 333-342.

Ventura, J., Wood, R., & Helleman, G. (2013). Symptom domains and neurocognitive functioning can help differentiate social cognitive processes in schizophrenia: A meta-analysis. *Schizophrenia Bulletin*, 39(1), 102-111.

Ventura, J., Subotnik, K. L., Ered, A., Gretchen-Doorly, D., Helleman, G. S., Vaskinn, A., & Nuechterlein, K. H. (2014). The relationship of attitudinal beliefs to negative symptoms, neurocognition, and daily functioning in recent-onset schizophrenia. *Schizophrenia Bulletin*, 40(6), 1308-1318.

Weijers, J., ten Kate, C., Eurelings-Bontekoe, E., Viechtbauer, W., Rampaart, R., Bateman, A., & Selten, J. P. (2016). Mentalization-based treatment for psychotic disorder: protocol of a randomized controlled trial. *BMC psychiatry*, 16(1), 191.

Williams, J., Stiles, M., & Shapiro, G. (1999). Cognitive Mechanisms in the Avoidance of Painful and Dangerous Thoughts: Elaborating the Assimilation Model. *Cognitive Therapy and Research*, 23(3), 285-306.

Wilson, J. S., & Costanzo, P. R. (1996). A preliminary study of attachment, attention, and schizotypy in early adulthood. *Journal of Social and Clinical Psychology, 15*(2), 231-260.